# SIMON W. YOUNG

MBChB FRACS MD Specialist Orthopaedic knee Surgeon, Associate Professor

## HIGH TIBIAL OSTEOTOMY (+/- ILIAC CREST BONE GRAFT HIP)

Post-Operative Care

## 1. WOUND CARE

After the surgery you will have a padded crepe bandage on your knee, you can remove these outer bandages 24-48 hours after the surgery.

There are plastic dressings under the crepe bandage, some blood staining on these is normal. If there is heavy blood staining or leaking you can contact us, or you can get your GP's practice nurse to change the dressings.

You can shower with these plastic dressings on, although you should not point the water straight at them or soak in a bath or pool.

The sutures used are absorbable and do not need to be removed. The dressings stay on until you are seen in clinic.

Icing of the knee is important following surgery, **especially in the first 72 hours**. Use a bag of crushed ice or frozen peas for up to 30 minutes, 4-5 times per day. After 72 hours ice use is optional.

# 2. ACTIVITY & WALKING

During the first few weeks ensure you get adequate rest and limit walking to what is necessary.

Elevate as much as possible and avoid long periods of time with the leg low down.

You will be restricted to 'Touch weight bearing' with crutches for the first four weeks. You can touch your foot to the ground, but do not put weight through the leg.

After 4 weeks you can put as much weight through the knee as is comfortable.

# 3. PHYSIOTHERAPY

Physiotherapy is an important part of your recovery, and usually starts six weeks following your surgery. Simon Young will discuss this with you.

### **FOLLOW UP**

Wendy will typically arrange a follow up appointment 2-3 weeks after the surgery, please liaise with her to ensure the date and time is suitable for you.

If you have any concerns prior to your appointment please phone 09 521 9849 or email ortho@axissportsmedicine.co.nz

After hours or if the query is urgent you can contact Simon Young on 021 616 183 or email simon.young@auckland.ac.nz

## HIGH TIBIAL OSTEOTOMY (+/- ILIAC CREST BONE GRAFT HIP)

**Questions & Answers** 

### 1. What medications do I take?

Your anaethetist will prescribe post operative pain medication. Local anaesthetic has been placed in your knee. This will typically wear off 6-8 hours after the surgery so you can expect pain to increase after this so ensure you begin taking the painkillers at the first sign pain is increasing.

#### 2. When can I drive?

Usually about 6 weeks, when you are completely off crutches.

## 3. When can I return to work or school?

This depends upon the demands of your job, but plan to take at least 14 days off of work or school. You can be back to a desk job within a week or two but physically demanding jobs may take 3 - 4 months to return to, once the osteotomy has healed.

## 4. When can I run?

Running is not allowed until the osteotomy has healed, your knee has a full range of motion, good strength and proprioception (balance) is attained. The earliest you can begin running straight ahead on a level surface is 4 months but often it takes 9-12 months. Your physiotherapist will help guide you.

### 5. Do I need a brace?

Not usually unless the osteotomy is complicated or there are additional procedures done.

If you have any concerns prior to your appointment please phone **09 521 9849** or email **ortho@axissportsmedicine.co.nz** 

After hours or if the query is urgent you can contact Simon Young on **021 616 183** or email **simon.young@auckland.ac.nz** 

